

Chapter 5: Relation

5.1 Relations and Their Properties

5.2 n-ary Relations and Their Applications

5.3 Representing Relations



Dr Patrick Chan

School of Computer Science and Engineering
South China University of Technology

Recall, Function is...

- Let **A** and **B** be nonempty sets
Function f from **A** to **B** is an **assignment** of **exactly one element of B** to **each element of A**
- By **defining** using a **relation**, a **function** from A to B contains **unique ordered pair (a, b)** for **every element $a \in A$**

AxB




Ch 5.1, 5.2, 5.3

Agenda

- What is Relation?
- Representation of Relation
 - Graph
 - Matrix
- Operators of Relation
- Properties of Relation

What is Relation?

Relation: x likes y



Ch 5.1, 5.2, 5.3

Relation

- Let A and B be sets
A **binary relation** from A to B is a **subset of A x B**
- Recall, for example:
 - $A = \{a_1, a_2\}$ and $B = \{b_1, b_2, b_3\}$
 - $A \times B = \{(a_1, b_1), (a_1, b_2), (a_1, b_3), (a_2, b_2), (a_1, b_2), (a_1, b_3)\}$

Relation: Example


- $S = \{\text{Peter, Paul, Mary}\}$
- $C = \{\text{C++}, \text{DisMath}\}$
- Given
 - Peter takes C++ Peter R C++ Peter $\not\text{R}$ DisMath
 - Paul takes DisMath Paul $\not\text{R}$ C++ Paul R DisMath
 - Mary takes none of them Mary $\not\text{R}$ C++ Mary $\not\text{R}$ DisMath
- $R = \{(\text{Peter, C++}), (\text{Paul, DisMath})\}$
- $(S \times C) - R = \text{ ~~}~~$

Relation

- R** is defined as
 - A **binary relation** from **A to B**
 - Ordered pairs**, which
 - First** element comes from **A**
 - Second** element comes from **B**
- aRb**: $(a, b) \in R$
- a~~R~~b**: $(a, b) \notin R$
- Moreover, when (a, b) belongs to R, **a** is said to be **related to b** by R

Relation VS Function


- Function** from a set A to a set B
 - All elements of A are assigned to B
 - Exactly one element of B to each element of A
- Relation** from a set A to a set B
 - Some elements of A are assigned to B
 - Zero, One or more elements of B to an element of A
- Function is a **special case** of Relation



Relation Representation Graph

- Relations can be represented by **Directed Graph**
 - You will learn the directed graph in detail in <Discrete Math Part 2>

- Graph $G = (V, E)$** consists of
 - a set of **vertices V**
 - a set of **edges E** ,
 - a connection between a pair of vertices




$V = \{ a, b, c, d \}$
 $E = \{ (a,b), (b,c), (b,d), (c,d) \}$

Relation Representation Graph

- G** to present a **relation** from **A** to **B** is


- vertices $V \subseteq A \cup B$
- edges $E \subseteq A \times B$




- For example
 - If there is an ordered pair (x, y) in R , then there is an edge from x to y in D



Relation Representation Graph




- Undirected Graph**
 - Edges are not directed
 - E.g. $(a, d) = (d, a)$



- Directed Graph (Digraph)**
 - Edges are directed
 - E.g. $(a, d) \neq (d, a)$

Relation Representation Graph: Example

- Peter R C++, Peter $\not R$ DisMath
 Paul $\not R$ C++, Paul R DisMath
 Mary $\not R$ C++, Mary $\not R$ DisMath




Directed Graph

Relation Representation Matrix

- Let **R** be a relation from $A = \{a_1, a_2, \dots, a_m\}$ to $B = \{b_1, b_2, \dots, b_n\}$
- An $m \times n$ connection **matrix M** for **R** is defined by

$$m_{ij} = \begin{cases} 1 & \text{if } (a_i, b_j) \in R \\ 0 & \text{if } (a_i, b_j) \notin R \end{cases}$$




	b_1	b_2	b_3	b_4
a_1	0	0	0	0
a_2	1	0	0	0
a_3	0	1	1	0
a_4	1	0	0	0
a_5	0	0	1	1

Relation on One Set

- Relation on the set **A** is a relation from **A** to **A**
 - Special case in relations
- Example:
 - $A = \{1, 2, 3, 4\}$
 - $R = \{(1,1), (1,4), (2,1), (2,3), (2,4), (3,1), (4,1), (4,2)\}$

Relation Representation Matrix: Example

- Peter R C++, Peter $\not R$ DisMath
- Paul $\not R$ C++, Paul R DisMath
- Mary $\not R$ C++, Mary $\not R$ DisMath




1	0
0	1
0	0

Directed Graph

Matrix

Relation on One Set Example 1

- Let **A** be the set $\{1, 2, 3, 4\}$, which ordered pairs are in the relation $R = \{(a, b) \mid a \text{ divides } b\}$?
- $R = \{(1,1), (1,2), (1,3), (1,4), (2,2), (2,4), (3,3), (4,4)\}$



1	1	1	1
0	1	0	1
0	0	1	0
0	0	0	1

Example 2

- How many different relations are there on a set with n elements?
- Suppose A has n elements
- Recall, a relation on a set A is a subset of $A \times A$
- $A \times A$ has n^2 elements
- If a set has m element, its has 2^m subsets
- Therefore, the answer is 2^{n^2}


Relation Properties

- Symmetric**
 $\forall a \forall b (((a, b) \in R) \rightarrow ((b, a) \in R))$
- Asymmetric** ((a,a) cannot be an element in R)
 $\forall a \forall b (((a, b) \in R) \rightarrow ((b, a) \notin R))$
- Antisymmetric** ((a,a) may be an element in R)
 $\forall a \forall b (((a, b) \in R \wedge (b, a) \in R) \rightarrow (a = b))$
- Asymmetry = Antisymmetry + Irreflexivity

Relation Properties

- Reflexive**
 $\forall a ((a, a) \in R)$
- Irreflexive**
 $\forall a ((a \in A) \rightarrow ((a, a) \notin R))$
- Transitive**
 $\forall a \forall b \forall c (((a,b) \in R \wedge (b,c) \in R) \rightarrow ((a,c) \in R))$


Relation Properties: Graph



Reflexive

$$\forall a ((a, a) \in R)$$


Every node has a self-loop



Irreflexive

$$\forall a ((a \in A) \rightarrow ((a, a) \notin R))$$

No node links to itself




Transitive


$$\forall a \forall b \forall c (((a,b) \in R \wedge (b,c) \in R) \rightarrow ((a,c) \in R))$$

Every two adjacent forms a triangle
(Not easy to observe in Graph)


Relation Properties: Graph



Symmetric
 $\forall a \forall b (((a, b) \in R) \rightarrow ((b, a) \in R))$
 Every link is bidirectional




Asymmetric
 $\forall a \forall b (((a, b) \in R) \rightarrow ((b, a) \notin R))$
 No link is bidirectional (Antisymmetric)
 No node links to itself (Irreflexive)




Antisymmetric
 $\forall a \forall b (((a, b) \in R \wedge (b, a) \in R) \rightarrow (a = b))$
 No link is bidirectional


Relation Properties: Matrix



Symmetric
 $\forall a \forall b (((a, b) \in R) \rightarrow ((b, a) \in R))$
 All identical across diagonal




Asymmetric
 $\forall a \forall b (((a, b) \in R) \rightarrow ((b, a) \notin R))$
 All 1's are across from 0's (Antisymmetric)
 All 0's on diagonal (Irreflexive)




Antisymmetric
 $\forall a \forall b (((a, b) \in R \wedge (b, a) \in R) \rightarrow (a = b))$
 All 1's are across from 0's

Relation Properties: Matrix



Reflexive
 $\forall a ((a, a) \in R)$
 All 1's on diagonal



Irreflexive
 $\forall a ((a \in A) \rightarrow ((a, a) \notin R))$
 All 0's on diagonal







Transitive
 $\forall a \forall b \forall c (((a, b) \in R \wedge (b, c) \in R) \rightarrow ((a, c) \in R))$
 Not easy to observe in Matrix

Example 1

- Consider the following relations on {1, 2, 3, 4}, Which properties these relations have?
 - $R_1 = \{(1,1), (1,2), (2,1), (2,2), (3,4), (4,1), (4,4)\}$
~~Reflexive~~ ~~Irreflexive~~ ~~Transitive~~ ~~Symmetric~~ ~~Asymmetric~~ ~~Antisymmetric~~
 - $R_2 = \{(1,1), (1,2), (2,1)\}$
~~Reflexive~~ ~~Irreflexive~~ ~~Transitive~~ ~~Symmetric~~ ~~Asymmetric~~ ~~Antisymmetric~~
 - $R_3 = \{(1,1), (1,2), (1,4), (2,1), (2,2), (3,3), (4,1), (4,4)\}$
~~Reflexive~~ ~~Irreflexive~~ ~~Transitive~~ ~~Symmetric~~ ~~Asymmetric~~ ~~Antisymmetric~~
 - $R_6 = \{(3,4)\}$
~~Reflexive~~ ~~Irreflexive~~ ~~Transitive~~ ~~Symmetric~~ ~~Asymmetric~~ ~~Antisymmetric~~

Example 2

	Reflexive Irreflexive Transitive Symmetric Asymmetric Antisymmetric
	Reflexive Irreflexive Transitive Symmetric Asymmetric Antisymmetric
	Reflexive Irreflexive Transitive Symmetric Asymmetric Antisymmetric
	Reflexive Irreflexive Transitive Symmetric Asymmetric Antisymmetric

Combining Relations

- As R is a subsets of $A \times B$, the set operations can be applied
 - Complement ($\bar{}$)
 - Union (\cup)
 - Intersection (\cap)
 - Difference ($-$)
 - Symmetric Complement (\oplus)

Example 3

- Let $A = \mathbb{Z}^+$, $R = \{ (a,b) \in A \times A \mid a \text{ divides } b \}$
 Is R symmetric, asymmetric, or antisymmetric?
- Symmetric** ($\forall a \forall b ((a, b) \in R \rightarrow (b, a) \in R)$) ❌
 - If aRb , it does not follow that bRa
- Asymmetric** ($\forall a \forall b ((a, b) \in R \rightarrow (b, a) \notin R)$) ❌
 - If $a=b$, then aRb and bRa
- Antisymmetric** ($\forall a \forall b ((a, b) \in R \wedge (b, a) \in R \rightarrow (a = b))$) ✓
 - If aRb and bRa , then $a=b$

Example

- Given, $A = \{1,2,3\}$, $B = \{1,2,3,4\}$
- $R_1 = \{(1,1), (2,2), (3,3)\}$,
 $R_2 = \{(1,1), (1,2), (1,3), (1,4)\}$
- $R_1 \cup R_2 = \{(1,1), (1,2), (1,3), (1,4), (2,2), (3,3)\}$
- $R_1 \cap R_2 = \{(1,1)\}$
- $R_1 - R_2 = \{(2,2), (3,3)\}$
- $R_2 - R_1 = \{(1,2), (1,3), (1,4)\}$
- $R_1 \oplus R_2 = \{(1,2), (1,3), (1,4), (2,2), (3,3)\}$

Combining Relations

- Let R be relation from a set A to a set B
- Inverse Relation** (R^{-1}) = $\{(b,a) \mid (a,b) \in R\}$
- Complementary Relation** (\overline{R}) = $\{(a,b) \mid (a,b) \notin R\}$
- Example**
 - $X = \{a, b, c\}$ $Y = \{1, 2\}$
 - $R = \{(a, 1), (b, 2), (c, 1)\}$
 - $R^{-1} = \{(1, a), (2, b), (1, c)\}$
 - $E = X \times Y = \{(a, 1), (b, 1), (c, 1), (a, 2), (b, 2), (c, 2)\}$
 - $\overline{R} = \{(a, 2), (b, 1), (c, 2)\} = E - R$

Ch 5.1, 5.2, 5.3

29

Combining Relations: Theorems

Example for the Proof

- Proof** $(R_1 \cup R_2)^{-1} = R_1^{-1} \cup R_2^{-1}$
- Assume**
 $(a,b) \in R_1$ & $(a,b) \in R_2$
- L.H.S.**
 - $(R_1 \cup R_2)^{-1} = \{(b,a) \mid (a,b) \in R_1 \vee (a,b) \in R_2\}$
 - $(R_1 \cup R_2)^{-1} = \{(b,a) \mid (a,b) \in R_1 \vee (a,b) \in R_2\}$
- R.H.S.**
 - $R_1^{-1} = \{(b,a) \mid (a,b) \in R_1\}$
 - $R_2^{-1} = \{(b,a) \mid (a,b) \in R_2\}$
 - $R_1^{-1} \cup R_2^{-1} = \{(b,a) \mid (a,b) \in R_1 \vee (a,b) \in R_2\}$

Recall...

- $A \cup B = \{x \mid x \in A \vee x \in B\}$
- $R^{-1} = \{(b,a) \mid (a,b) \in R\}$

Ch 5.1, 5.2, 5.3

31

Combining Relations Theorems

- Let R_1 and R_2 be relations from A to B . Then
 - $(R^{-1})^{-1} = R$
 - $(R_1 \cup R_2)^{-1} = R_1^{-1} \cup R_2^{-1}$
 - $(R_1 \cap R_2)^{-1} = R_1^{-1} \cap R_2^{-1}$
 - $(A \times B)^{-1} = B \times A$
 - $\emptyset^{-1} = \emptyset$
 - $(\overline{R})^{-1} = \overline{(R^{-1})}$
 - $(R_1 - R_2)^{-1} = R_1^{-1} - R_2^{-1}$
 - If $R_1 \subseteq R_2$ then $R_1^{-1} \subseteq R_2^{-1}$

Ch 5.1, 5.2, 5.3

30

Combining Relations Example 1

- Given
 - R_1 is **symmetric**
 - R_2 is **antisymmetric**
- Does it $R_1 \cup R_2$ is **transitive**?
- Not transitive** by giving a **counterexample**
 - $R_1 = \{(1,2), (2,1)\}$ which is **symmetric**
 - $R_2 = \{(1,2), (1,3)\}$ which is **antisymmetric**
 - $R_1 \cup R_2 = \{(1,2), (2,1), (1,3)\}$, **not transitive**

Ch 5.1, 5.2, 5.3

32

Example 2

- Given R_1 and R_2 are transitive on A
- Does $R_1 \cup R_2$ is transitive?
- Not transitive by giving a counterexample
 - $A = \{1, 2\}$
 - $R_1 = \{(1,2)\}$, which is transitive
 - $R_2 = \{(2,1)\}$, which is transitive
 - $R_1 \cup R_2 = \{(1,2), (2,1)\}$, not transitive

Combining Relations: Matrix

- Example

$$M_{R_1} = \begin{bmatrix} 1 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix} \quad M_{R_1 \cup R_2} = M_{R_1} \vee M_{R_2} = \begin{bmatrix} 1 & 0 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 0 \end{bmatrix}$$

$$M_{R_2} = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 0 & 0 \end{bmatrix} \quad M_{R_1 \cap R_2} = M_{R_1} \wedge M_{R_2} = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

Combining Relations: Matrix

- Suppose that R_1 and R_2 are relations on a set A represented by the matrices M_{R_1} and M_{R_2} , respectively

- Join operator (OR)


$$M_{R_1 \cup R_2} = M_{R_1} \vee M_{R_2}$$

- Meet operator (AND)

$$M_{R_1 \cap R_2} = M_{R_1} \wedge M_{R_2}$$

Combining Relations Composite

- Recall, the composition in functions...
- Let
 - g be a function from the set A to the set B
 - f be a function from the set B to the set C
- The composition of the functions f and g , denoted by $f \circ g$, is defined by $(f \circ g)(a) = f(g(a))$





Combining Relations Composite

- Let
 - R be a relation from a set A to a set B
 - S be a relation from a set B to a set C
- The **composite** of R and S is the relation consisting of ordered pairs (a, c), where
 - a ∈ A, c ∈ C, and
 - There **exists** an element b ∈ B, such that (a, b) ∈ R and (b, c) ∈ S
- The composite of R and S is denoted by **S o R**


Combining Relations Composite: Example

- What is the composite of the relations R and S, where
 - R is the relation from {1,2,3} to {1,2,3,4} with $R = \{(1,1), (1,4), (2,3), (3,1), (3,4)\}$
 - S is the relation from {1,2,3,4} to {0,1,2} with $S = \{(1,0), (1,2), (2,0), (3,1), (3,2), (4,1)\}$?
- $S \circ R = \{(1,0), (1,2), (1,1), (2,1), (2,2), (3,0), (3,2), (3,1)\}$



Combining Relations Composite

- Suppose
 - R be a relation from a set A to a set B
 - S be a relation from a set B to a set C
- $(x, y) \in S \circ R$ implies $\exists z ((x, z) \in R \wedge (z, y) \in S)$**



Remark: May be more than one element z, where (x, z) ∈ R and (z, y) ∈ S

Combining Relations Composite: Properties

- Let R_1 and R_2 be relations on the set A.
- Show $(R_1 \circ R_2)^{-1} = R_2^{-1} \circ R_1^{-1}$
- Proof:

Let $(x, y) \in (R_1 \circ R_2)^{-1}$

$$\begin{aligned}
 (x, y) &\in (R_1 \circ R_2)^{-1} \\
 \Leftrightarrow (y, x) &\in R_1 \circ R_2 \\
 \Leftrightarrow \exists z ((y, z) \in R_2 \wedge (z, x) \in R_1) \\
 \Leftrightarrow \exists z ((z, y) \in R_2^{-1} \wedge (x, z) \in R_1^{-1}) \\
 \Leftrightarrow (x, y) &\in R_2^{-1} \circ R_1^{-1}
 \end{aligned}$$


$(x, y) \in S \circ R$ implies $\exists z ((x, z) \in R \wedge (z, y) \in S)$

Composite: Properties

- Let F, G and H be relations on the set A , then
 - $F \circ (G \cup H) = (F \circ G) \cup (F \circ H)$
 - $F \circ (G \cap H) \subseteq (F \circ G) \cap (F \circ H)$
 - $(G \cup H) \circ F = (G \circ F) \cup (H \circ F)$
 - $(G \cap H) \circ F \subseteq (G \circ F) \cap (H \circ F)$

Composite: Example

- Let $R = \{(1,1), (2,1), (3,2), (4,3)\}$
- Find the powers $R^n, n = 2,3,4,\dots$
- $R^2 = R \circ R = \{(1,1), (2,1), (3,1), (4,2)\}$
- $R^3 = R^2 \circ R = \{(1,1), (2,1), (3,1), (4,1)\}$
- $R^4 = R^3 \circ R = \{(1,1), (2,1), (3,1), (4,1)\}$
- $R^n = R^3$ for $n = 5, 6, 7, \dots$



Composite

- Let R be a relation on the set A . The powers $R^n, n = 1, 2, 3, \dots$, are defined recursively by
 - $R^1 = R$
 - $R^2 = R \circ R$
 - $R^3 = R^2 \circ R = (R \circ R) \circ R$
 - ...
 - $R^{n+1} = R^n \circ R$

Composite: Matrix

- Suppose
 - R_1 be relation from set A to set B represented by M_{R_1}
 - R_2 be relation from set B to set C represented by M_{R_2}
- The matrix for the composite of R_1 and R_2 is:

$$M_{R_2 \circ R_1}$$
 - Size of M_{R_1} and M_{R_2} is $|A| \times |B|$ and $|B| \times |C|$
 - Size of $M_{R_2 \circ R_1}$ is $|A| \times |C|$

Composite: Matrix

Define:

n : the number of row of R_1
the number of column of R_2

$$M_{R_2 \circ R_1} = M_{R_2} \odot M_{R_1}$$

where $(M_{R_2} \odot M_{R_1})_{ij} = \bigvee_{k=1}^n [(M_{R_1})_{ik} \wedge (M_{R_2})_{kj}]$


An element in the matrix

Such that

$$(M_{R_2} \odot M_{R_1})_{ij} = 1$$

if and only if

$$(M_{R_1})_{ik} = (M_{R_2})_{kj} = 1 \text{ for some } k$$



Composite: Matrix

The powers R^n can be defined using matrix as:

$$M_{R^n} = (M_R)^n$$

Example

Find the matrix representing the relation R^2

$$M_R = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 1 & 1 \\ 1 & 0 & 0 \end{bmatrix}$$

$$M_{R^2} = (M_R)^2 = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 1 & 1 \\ 1 & 0 & 0 \end{bmatrix} \odot \begin{bmatrix} 0 & 1 & 0 \\ 0 & 1 & 1 \\ 1 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 1 & 1 \\ 1 & 1 & 1 \\ 0 & 1 & 0 \end{bmatrix}$$

Composite: Matrix: Example

$$M_R = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

3 x 4

n : the number of column of R_1
the number of row of R_2

$$M_S = \begin{bmatrix} 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \end{bmatrix}$$

4 x 5

$$M_{S \circ R} = \begin{bmatrix} 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 0 \end{bmatrix}$$

$i = 1, j = 1$
 $k = 1$

$n=4$

$$(M_{R_2} \odot M_{R_1})_{ij} = \bigvee_{k=1}^n [(M_{R_1})_{ik} \wedge (M_{R_2})_{kj}]$$

Composite: Property 1


Theorem

If $R \subset S$, then $S \circ R \subset S \circ S$

- Assume $(x,y) \in S \circ R$, there exists an element z , which $(x,z) \in R$ and $(z,y) \in S$
- As $R \subset S$ and $(x,z) \in R$, $(x,z) \in S$
- Therefore, as $(x,z) \in S$ and $(z,y) \in S$, $(x,y) \in S \circ S$
- $S \circ R \subset S \circ S$
- It implies:
If $R \subset S$ and $T \subset U$, then $R \circ T \subset S \circ U$

Composite: Property 2

An ordered pairs (x, y) is in R^n iff there is a path of length n from x to y in R




Composite: Property 2

Theorem

Let R be a relation on A . There is a path of length n from a to b in R iff $(a, b) \in R^n$

Composite: Property 2

An ordered pairs (x, y) is in R^n iff there is a path of length n from x to y in R



Example

- In R , $1 > 2 > 4$, length = 2 $\Leftrightarrow (1,4) \in R^2$
- In R , $3 > 2 > 4 > 3$, length = 3 $\Leftrightarrow (3,3) \in R^3$
- $(1,2) \in R^4 \Leftrightarrow$ In R , $1 > 2 > 4 > 3 > 2$, length = 4

Composite: Property 2

Proof by Induction

a path of length n from a to b iff $(a, b) \in R^n$

- Show $n=1$ is true
 - An arc from a to b is a path of length 1, which is in $R^1 = R$
 - Hence the assertion is true for $n = 1$
- Assume it is true for k . Show it is true for $k+1$
 - As it is true for $n = 1$, suppose (a, x) is a path of length 1, then $(a, x) \in R$
 - As it is true for $n = k$, suppose (x, b) is a path of length k , then $(x, b) \in R^k$
 - Considering, $(a, x) \in R$ and $(x, b) \in R^k$, $(a, b) \in R^{k+1} = R^k \circ R$ as there exists an element x , such that $(a, x) \in R$ and $(x, b) \in R^k$
 - The length of (a,b) is $k+1$

Composite: Property 3

- R is transitive iff $R^n \subseteq R$ for $n > 0$.

- Proof

1. $(R^n \subseteq R) \rightarrow R$ is transitive

- Suppose $(a,b) \in R$ and $(b,c) \in R$
- (a,c) is an element of R^2 as $R^2 = R \circ R$
- As $R^2 \subseteq R$, $(a,c) \in R$
- Hence R is transitive

Composite: Property 4

- Proof: If R is transitive, R^n is also transitive

- When $n = 1$, R is transitive

- Assume R^k is transitive

- Show R^{k+1} is transitive

Given $(a,b) \in R^{k+1}$ and $(b,c) \in R^{k+1}$, show $(a,c) \in R^{k+1}$

- $R^{k+1} = R^k \circ R$
- As $(a,b) \in R^{k+1}$, $(d,b) \in R^k$ and $(a,d) \in R$
- As $(b,c) \in R^{k+1}$, $(f,c) \in R^k$ and $(b,f) \in R$
- As $(a,c) \in R^{k+1}$, $(?,c) \in R^k$ and $(a,?) \in R$

Composite: Property 3**2. R is transitive $\rightarrow (R^n \subseteq R)$**

- Use a proof by induction:

- **Basis:** Obviously true for $n = 1$.

- **Induction:** Assume true for n , show it is true for $n + 1$

- For any (x, y) is in R^{n+1} , there is a z such that $(x, z) \in R$ and $(z, y) \in R^n$
- But since $R^n \subseteq R$, $(z, y) \in R$
- As R is transitive, (x, z) and (z, y) are in R , so (x, y) is in R
- Therefore, $R^{n+1} \subseteq R$

Composite: Property 4

- Given $(a,b) \in R^{k+1}$ and $(b,c) \in R^{k+1}$, show $(a,c) \in R^{k+1}$

- $R^{k+1} = R^k \circ R$
- As $(a,b) \in R^{k+1}$, $(d,b) \in R^k$ and $(a,d) \in R$
- As $(b,c) \in R^{k+1}$, $(f,c) \in R^k$ and $(b,f) \in R$
- As $(a,c) \in R^{k+1}$, $(?,c) \in R^k$ and $(a,?) \in R$

- As “ R is transitive iff $R^n \subseteq R$ for $n > 0$ ”

- $(d,b) \in R^k \subseteq R$

- As R is transitive, $(d,b) \in R$ and $(b,f) \in R$ imply $(d,f) \in R$

- As R is transitive, $(d,f) \in R$ and $(a,d) \in R$ imply $(a,f) \in R$

- Therefore, by considering, $(f,c) \in R^k$ and $(a,f) \in R$, $(a,c) \in R^{k+1}$

Composite: Property 4

- Proof: If R is transitive, R^n is also transitive
- When $n = 1$, R is transitive
- Assume R^k is transitive
- Show R^{k+1} is transitive

Given $(a,b) \in R^{k+1}$ and $(b,c) \in R^{k+1}$, show $(a,c) \in R^{k+1}$

- $R^{k+1} = R^k \circ R$
- As $(a,b) \in R^{k+1}$, $(a,d) \in R^k$ and $(d,b) \in R$
- As $(b,c) \in R^{k+1}$, $(b,f) \in R^k$ and $(f,c) \in R$
- As $(a,c) \in R^{k+1}$, $(a,?) \in R^k$ and $(?,c) \in R$

n-ary Relation

- Let A_1, A_2, \dots, A_n be sets
An **n-ary relation** on these sets is a subset of $A_1 \times A_2 \times \dots \times A_n$
- **Domains** of the relation: the sets A_1, A_2, \dots, A_n
- **Degree** of the relation: n

Composite: Property 4

- Given $(a,b) \in R^{k+1}$ and $(b,c) \in R^{k+1}$, show $(a,c) \in R^{k+1}$
 - $R^{k+1} = R^k \circ R$
 - As $(a,b) \in R^{k+1}$, $(a,d) \in R^k$ and $(d,b) \in R$
 - As $(b,c) \in R^{k+1}$, $(b,f) \in R^k$ and $(f,c) \in R$
 - As $(a,c) \in R^{k+1}$, $(a,?) \in R^k$ and $(?,c) \in R$
- As “ R is transitive iff $R^n \subseteq R$ for $n > 0$ ”
- $(b,f) \in R^k \subseteq R$
- As R is transitive, $(d,b) \in R$ and $(b,f) \in R$ imply $(d,f) \in R$
- As R is transitive, $(d,f) \in R$ and $(f,c) \in R$ imply $(d,c) \in R$
- Therefore, by considering, $(a,d) \in R^k$ and $(d,c) \in R$, $(a,c) \in R^{k+1}$

n-ary Relation: Example

- Let R be the relation on $Z \times Z \times Z^+$ consisting of triples (a, b, m) , where a, b , and m are integers with $m \geq 1$ and $a = b \pmod{m}$, (i.e. m divides $a-b$)
- Degree of the relation? **3**
- First domain is: **the set of all integers**
- Second domain is: **the set of all integers**
- Third domain: **the set of positive integers**
- Do they belong to R ?

■ $(8,2,3)$	Y	■ $(7,2,3)$	N
■ $(-1,9,5)$	Y	■ $(-2,-8,5)$	N

Relational Database VS n-ary Relation

- A **database** consists of **records** made up of **fields**

- Each **record** is a **n-tuple** (n fields)

- For example:

ID num	Name	Major	GPA
888323	Adams	Data Structure	85
231455	Peter	C++	61

- **Domain:** ID num, Name, Major, GPA

- **Relation:** (888323, Adams, Data Structure, 85),
(231455, Sam, C++, 61)

- Relations are displayed as tables

ID_number	Student_name	Major	Grade
888323	Adams	Data Structure	85
231455	Peter	C++	61
678543	Sam	Data Structure	98

Relational Database VS n-ary Relation

- n-ary relation can be:

- **Determining** all n-tuples **satisfy** certain **conditions**

- **Joining** the records in **different tables**

ID_number	Major	Grade
888323	Data Structure	85
231455	C++	61
678543	Data Structure	98
453876	Discrete Math	83

ID_number	Student_name
231455	Adams
888323	Peter
102147	Sam
453876	Goodfriend
678543	Rao
786576	Stevens