

Chapter 9: Graphs

9.1

Graphs and Graph Model

9.2

Graph Terminology and Special Types of Graphs

9.3

Representing Graphs and graph Isomorphism

9.4

Connectivity

Dr Patrick Chan

School of Computer Science and Engineering
South China University of Technology


Agenda

- Graph
- Terminology
- Connectivity
- Isomorphism

2

Graph

- A graph $G = (V, E)$ consists of a set of vertices V , and a set of edges E


 $V = \{ a, b, c, d, e, f \}$
 $E = \{ (a,b), (a,c), (a,d), (b,e), (c,d), (c,e), (d,e), (e,f) \}$

- Vertices (V)
 - $|V|$: the number of vertices
- Edges (E)
 - Sometimes referred as **arc**
 - Connection between a pair of vertices (v, w) , where v and w belong to V
 - $|E|$: the number of edges
 - **Weight** may be included

3


Graph Structure

- Key questions about Graph Structure
 - Directed / Undirected Edge?
 - Single / Multiple Connection?
 - Loop?

4


Directed/Undirected?

- **Undirected Graph**
 - Edges are **not directed**
 - If (a,b), then (b,a)
- **Directed Graph (Digraph)**
 - Edges are **directed**
 - (a,b) does not mean (b,a)




Single/Multiple Connection?

- **Simple Graph**
 - No two **edges connects** the same **pair of vertices**
 - Loop is not allowed
- **Multigraph**
 - Two **vertices** may be **connected by more than one edges**
 - An edge cannot be identified uniquely by a pair of vertices
 - **Additional name is needed**
 - E.g. (c,d) means u or v



Loop?

- **Multigraph** does **not allow loop**
- **Pseudograph** is a special multigraph **allows loop**
- Sometimes, the meanings of **Pseudograph** and **Multigraph** are the **same**



Summary

		No Loop	Loop
Undirected	Single Edge	Simple Graph	/
	Multiple Edge	Multigraph	Pseudograph (Multigraph)

		No Loop	Loop
Directed	Single Edge	Simple Directed Graph	/
	Multiple Edge	Directed Multigraph	Mixed Graph

Adjacent / Neighbor

Undirected graph

- Let (v_1, v_2) is an edge
- v_1 and v_2 are endpoints
- v_1 is adjacent to v_2
- Also mean
“ v_2 is adjacent to v_1 ”
since $(v_1, v_2) = (v_2, v_1)$



Directed graph


- Let (v_1, v_2) is an edge
- v_1 is initial vertex
- v_2 is terminal (end) vertex
- v_1 is adjacent to v_2
- v_2 is adjacent from v_1
- Do not mean
“ v_2 is adjacent to v_1 ”



9


Adjacent / Neighbor

- e incidents with v_1 and v_2
- e connects v_1 and v_2




Example:

a & b are adjacent
b & a are adjacent





w is adjacent to z
z is not adjacent to w



10

Neighbor Set

- Neighbor Set $N(v)$ contains all adjacent vertices of v
- For example: $N(c) = \{a, b, d\}$
 $N(z) = \{y\}$




11

Degree


Undirected graph

- Degree:** number of edges containing that vertex (Adjacent vertex number)
- Isolated vertex:** $\text{deg} = 0$
- Pendant vertex:** $\text{deg} = 1$
- E.g. $\text{deg}(c) = 3$



Directed graph

- In-Degree:** in-bound edge number
- Out-Degree:** out-bound edge number
- E.g. $\text{deg}^-(z) = 2$
 $\text{deg}^+(z) = 1$




12

Degree: Example


What are the degrees of the following vertices?

- deg(a) = 2 deg(g) = 2 deg+(g) = 0
- deg(b) = 3 deg(h) = 0 deg+(h) = 3
- deg(c) = 2 deg(i) = 1 deg+(i) = 1
- deg(d) = 2 deg(j) = 1 deg+(j) = 1
- deg(e) = 0 deg(k) = 0 deg+(k) = 0
- deg(f) = 1 deg(l) = 1 deg+(l) = 0



Degree Sequence

A degree sequence is a monotonic nonincreasing sequence of the degrees of vertices in an undirected graph.



- (2,3,2,2,0,1) Not monotonic nonincreasing
- (3,2,2,2,1,0) Degree sequence
- (3,3,2,2,2) Degree sequence


Degree Handshaking Theorem 1

For any undirected graph $G = (V, E)$,

$$2|E| = \sum_{v \in V} \text{deg}(v)$$

Twice number of edges = sum of degrees

- Each edge maps to two vertices (start & end)
- It also applies to multiple edges and loop



Degree: Handshaking Theorem 1 Example 1

How many edges are there in a graph with 10 vertices each of degree six?

- Total degree = $10 \times 6 = 60$
- According to Handshaking Theorem


$$2|E| = \sum_{v \in V} \text{deg}(v)$$

$|E| = 60/2 = 30$


Example 1

- Is there a graph with degree sequence...


- (2,2,2)? **Yes**




- (3,3,3,3)? **Yes**



- (2,2,1)? **No, not even**




- (5,5,4)? **Yes**




Handshaking Theorem 2


- Undirected graph has an even number of vertices of odd degree



vertices of odd degree
2



vertices of odd degree
4



vertices of odd degree
2

Handshaking Theorem 2

Proof

- Let V_o and V_e be the set of vertices of odd and even degree

$$2|E| = \underbrace{\sum_{v \in V} \deg(v)}_{\text{even}} = \underbrace{\sum_{v \in V_o} \deg(v)}_{\text{also be even}} + \underbrace{\sum_{v \in V_e} \deg(v)}_{\text{Must be even}}$$


- As summation of even degree (2nd term) is even
- Summation of odd degree (1st term) is also even
 - As $\deg(v)$ is odd for $v \in V_o$
 - The number of $\deg(v)$ must be even for $v \in V_o$

Handshaking Theorem 3

- For any directed graph $G = (V, E)$,

$$|E| = \sum_{v \in V} \deg^+(v) = \sum_{v \in V} \deg^-(v)$$

- Each edge maps to one initial and on end vertices




$|E| = 4$

$\deg^+(a) = 2$ $\deg^-(a) = 0$
 $\deg^+(b) = 0$ $\deg^-(b) = 3$
 $\deg^+(c) = 1$ $\deg^-(c) = 1$
 $\deg^+(d) = 1$ $\deg^-(d) = 0$

- It also applies to multiple edges and loop

Path

- A sequence of vertices v_1, v_2, \dots, v_n of length $n-1$ with an edge from v_i to v_{i+1} for $1 \leq i < n$
- A path is **simple** if all vertices on the path are distinct




Vertex a to e
 $a > b > e$ Length = 2 Simple
 $a > c > b > e$ Length = 3 Simple

Vertex a to b
 $a > b$ Length = 1 Simple
 $a > c > b$ Length = 2 Simple
 $a > c > \textcircled{b} > e > d > c > \textcircled{b}$ Length = 6 Not Simple

Cycle (Circuit)


- A path connects v_i to itself
- A cycle is **simple** if the path is simple, except the first and last vertices are the same



$a > c > b > a$ Simple Cycle
 $b > e > d > c > b$ Simple Cycle
 $\textcircled{b} > e > d > c > \textcircled{b} > a > c > \textcircled{b}$ Cycle


Acyclic

- A graph without cycle is called **acyclic**
- A directed graph without cycles is called a **Directed Acyclic Graph (DAG)**



Connectedness

- Vertices v, w are connected if and only if there is a path starting at v and ending at w
- Every graph consists of **separate connected pieces** called **connected components**




Are a and e connected? Yes
 Are a and d connected? No

How many connected components?
 2


Connectedness

- Undirected graph**
 - Connected:** if there is at least one path from any vertex to any other (Only one connected component)
- Directed graph**
 - Weakly connected:** Directed graph without considering directions is connected
 - Strongly connected:** Directed graph with considering direction is connected



Maximal/Minimal graph


- A graph G is said to be a **maximal graph** (**minimal graph**) with respect to a **property P** if G has property P and no proper **supergraph** (**subgraph**) of G has the property P



Graph Operation



- Given $G_1=(V_1,E_1)$ and $G_2=(V_2,E_2)$
 - Complement \bar{G}_1**
($V_1, \{uv \mid u \neq v, uv \notin E_1\}$)
 - Intersection $G_1 \cap G_2$**
($V_1 \cap V_2, E_1 \cap E_2$)
 - Union $G_1 \cup G_2$**
($V_1 \cup V_2, E_1 \cup E_2$)
 - Join $G_1 + G_2$** if $V_1 \cap V_2 = \emptyset$
($V_1 \cup V_2, E_1 \cup E_2 \cup \{uv \mid u \in V_1 \text{ and } v \in V_2\}$)
 - Cartesian Product $G_1 \times G_2$**
($V_1 \times V_2, \{(u_1, u_2), (v_1, v_2) \mid (u_1 = v_1 \text{ and } \{u_2, v_2\} \in E_2) \text{ or } (u_2 = v_2 \text{ and } \{u_1, v_1\} \in E_1)\}$)

Graph Operation Example 1



- Complement \bar{G}_1**
($V_1, \{uv \mid u \neq v, uv \notin E_1\}$)
- Intersection $G_1 \cap G_2$**
($V_1 \cap V_2, E_1 \cap E_2$)
- Union $G_1 \cup G_2$**
($V_1 \cup V_2, E_1 \cup E_2$)
- Join $G_1 + G_2$** if $V_1 \cap V_2 = \emptyset$
($V_1 \cup V_2, E_1 \cup E_2 \cup \{uv \mid u \in V_1 \text{ and } v \in V_2\}$)
- Cartesian Product $G_1 \times G_2$**
($V_1 \times V_2, \{(u_1, u_2), (v_1, v_2) \mid (u_1 = v_1 \text{ and } \{u_2, v_2\} \in E_2) \text{ or } (u_2 = v_2 \text{ and } \{u_1, v_1\} \in E_1)\}$)


Graph Operation Example 2



- **Complement \bar{G}_1**
($V_1, \{uv \mid u \neq v, uv \notin E_1\}$)
- **Intersection $G_1 \cap G_2$**
($V_1 \cap V_2, E_1 \cap E_2$)
- **Union $G_1 \cup G_2$**
($V_1 \cup V_2, E_1 \cup E_2$)
- **Join $G_1 + G_2$ if $V_1 \cap V_2 = \emptyset$**
($V_1 \cup V_2, E_1 \cup E_2 \cup \{uv \mid u \in V_1 \text{ and } v \in V_2\}$)
- **Cartesian Product $G_1 \times G_2$**
($V_1 \times V_2, \{(u_1, u_2), (v_1, v_2) \mid (u_1 = v_1 \text{ and } \{u_2, v_2\} \in E_2) \text{ or } (u_2 = v_2 \text{ and } \{u_1, v_1\} \in E_1)\}$)


Graph Operation Example 3

- **Complement \bar{G}_1**
($V_1, \{uv \mid u \neq v, uv \notin E_1\}$)
- **Intersection $G_1 \cap G_2$**
($V_1 \cap V_2, E_1 \cap E_2$)
- **Union $G_1 \cup G_2$**
($V_1 \cup V_2, E_1 \cup E_2$)
- **Join $G_1 + G_2$ if $V_1 \cap V_2 = \emptyset$**
($V_1 \cup V_2, E_1 \cup E_2 \cup \{uv \mid u \in V_1 \text{ and } v \in V_2\}$)
- **Cartesian Product $G_1 \times G_2$**
($V_1 \times V_2, \{(u_1, u_2), (v_1, v_2) \mid (u_1 = v_1 \text{ and } \{u_2, v_2\} \in E_2) \text{ or } (u_2 = v_2 \text{ and } \{u_1, v_1\} \in E_1)\}$)




Graph Operation Example 4

- **Complement \bar{G}_1**
($V_1, \{uv \mid u \neq v, uv \notin E_1\}$)
- **Intersection $G_1 \cap G_2$**
($V_1 \cap V_2, E_1 \cap E_2$)
- **Union $G_1 \cup G_2$**
($V_1 \cup V_2, E_1 \cup E_2$)
- **Join $G_1 + G_2$ if $V_1 \cap V_2 = \emptyset$**
($V_1 \cup V_2, E_1 \cup E_2 \cup \{uv \mid u \in V_1 \text{ and } v \in V_2\}$)
- **Cartesian Product $G_1 \times G_2$**
($V_1 \times V_2, \{(u_1, u_2), (v_1, v_2) \mid (u_1 = v_1 \text{ and } \{u_2, v_2\} \in E_2) \text{ or } (u_2 = v_2 \text{ and } \{u_1, v_1\} \in E_1)\}$)




$K_2 + K_3$




$\bar{K}_2 + \bar{K}_3$

Graph Operation Example 5

- **Complement \bar{G}_1**
($V_1, \{uv \mid u \neq v, uv \notin E_1\}$)
- **Intersection $G_1 \cap G_2$**
($V_1 \cap V_2, E_1 \cap E_2$)
- **Union $G_1 \cup G_2$**
($V_1 \cup V_2, E_1 \cup E_2$)
- **Join $G_1 + G_2$ if $V_1 \cap V_2 = \emptyset$**
($V_1 \cup V_2, E_1 \cup E_2 \cup \{uv \mid u \in V_1 \text{ and } v \in V_2\}$)
- **Cartesian Product $G_1 \times G_2$**
($V_1 \times V_2, \{(u_1, u_2), (v_1, v_2) \mid (u_1 = v_1 \text{ and } \{u_2, v_2\} \in E_2) \text{ or } (u_2 = v_2 \text{ and } \{u_1, v_1\} \in E_1)\}$)




$\bar{K}_2 \times K_3$




$K_2 \times \bar{K}_3$

Complete Graph

- Complete graph K_n if there is an edge between every pair of vertices, where n is the number of vertices
- Complete Undirected Graph
- Complete Directed Graph



K_4




Cycle Graph


- Cycle graph C_n is a circular graph with $V = \{0, 1, 2, \dots, n-1\}$ where vertex i is connected to $(i+1) \bmod n$ and to $(i-1) \bmod n$
 - like a polygon




C_1



C_2




C_3




C_4

Wheel Graph


- Wheel graph W_n is a cycle graph with an extra vertex in the middle which contact to each of other vertices




W_1



W_2



W_3



W_4

Cube


- n -cube Q_n is defined recursively.
 - Q_0 is just a vertex
 - Q_{n+1} is gotten by taking 2 copies of Q_n and joining each vertex v of Q_n with its copy v'




Q_0




Q_1



Q_2



Q_3




Q_4

Type of Graph

Bipartite Graph

- A graph is **bipartite** if **all vertices** can be **separated into two partitions**, (i.e. $V = V_1 \cup V_2$ and $\emptyset = V_1 \cap V_2$) so that any two adjacent vertices are in different partitions
 - (V_1, V_2) is called a bipartition of V of G




37

Type of Graph

Bipartite Graph: Theorem

- A simple graph is **bipartite if and only** if it is possible to assign one of **two different colors** to each vertex of the graph so that **no two adjacent vertices are assigned the same color**




38

Type of Graph

Bipartite Graph: Example 1

- Is C_n (**Cycle graph**) bipartite?
 - When n is **even**, **Yes**
 - All odd vertices are in a color and all vertices numbers are in another color
 - All vertices are only adjacent to opposite color
 - When n is **odd**, **No** (except $n = 1$)
 - Both n and 1 are odd, but n^{th} vertex is next to the 1^{st} vertex




39

Type of Graph

Bipartite Graph: Example 2



- Is the given graph **bipartite**?
 - NO**
 - For example, consider a , b , and c . There is two adjacent vertices are assigned the same color if only two colors are allowed



40


Complete Bipartite Graph

- When **all possible edges exist in a simple bipartite graph** with **m and n** vertices in two **partitions**, the graph is called **complete bipartite $K_{m,n}$**




Subgraph

- Let $G = (V,E)$ and $H = (W,F)$ be graphs. H is a **subgraph** of G , if $W \subseteq V$ and $F \subseteq E$
 - Subgraph is a graph inside another group
- A **subgraph** H of G is a **proper subgraph** of G if $H \neq G$





Subgraph: Example




- How many different Q_2 subgraphs does Q_3 have?

6




Tree

- Tree** is an **undirected, connected and acyclic** graph
 - n vertices has $(n-1)$ edges




- Forest** is an **undirected, disconnected, acyclic** graph
 - Disjoint collection of trees



Theorem 1

- A **tree** with at least two vertices has **at least two leaves**
- Assume P is a **longest path** in a tree T
- **Prove** its endpoints are leaves
- Suppose v is **not a leaf**, then v has **at least two neighbors**, x and y
- One of them (say x) must **not in P** , otherwise a cycle
- Let P' be the **path** that **begins at x** followed by P
- This is a **longer path than P** which is **contradict to the assumption**



45

Theorem 2

- A **tree** on n vertices has **$n - 1$ edges**
- For $N(1)$
 - If $n = 1$, then T has no edges
- **Assume $N(k)$ is true**
 - T with n vertices has exactly $n - 1$
- **Show $N(k+1)$**

46

Theorem 2

- Show T with $n+1$ vertices has **exactly $n - 1$**
- Since T is a tree, T has **at least two leaves (Theorem 1)**
- Let T' be the graph created by **deleting a leaf in T**
- Note that T' is a **tree with n vertices**, since:
 - T' is **connected** and **acyclic**
 - T' has **one less vertex** than T
- According to $N(k)$, T' has **$n - 1$ edges**
- Since T' has one less edge than T , T has **K edges**

47



Theorem 3

- Let G be a graph with n vertices. Then the following are equivalent:
 1. G is a **tree**
 2. G is a **maximal acyclic graph**
 3. G is a **minimal connected graph**
 4. G is **acyclic** and it has **$n - 1$ edges**
 5. G is **connected** and it has **$n - 1$ edges**
 6. Between any **two distinct vertices** of G there exists a **unique path**


48

Spanning Tree


- Spanning Tree in a connected graph G is a sub-graph H of G that includes all the vertices of G and is also a tree




Not Spanning Tree
(not all vertices)



Spanning Tree





Not Spanning Tree
(not a tree)




Spanning Tree

Minimum Spanning Tree


- Minimum Spanning Trees (MST) is a spanning tree with the minimal cost to call all the vertices




Not Spanning Tree
(not all vertices)



Not Spanning Tree
(not a tree)





Spanning Tree
MST (22)




Spanning Tree
Not MST (31)

Graph Isomorphism


- Is the following graphs the same as Graph A?



Yes
Different Labels



Yes
Different Positions




Yes
Different Label
and Positions

Graph Isomorphism

- Two graphs, A and B , which contain the same number of graph vertices connected in the same way are said to be **isomorphic**, $A \cong B$
- Applications
 - Checking fingerprint
 - Testing molecules

Graph Isomorphism

- If $G_1 \cong G_2$, do they have
 - the same number of vertices? **Yes**
 - the same number of edges? **Yes**
 - the same degree sequence? **Yes**
- Are $G_1 \cong G_2$, if they have
 - the same number of vertices? **No**
 - the same number of edges? **No**
 - the same degree sequence? **No**




Graph Isomorphism

- $G_1 = (V_1, E_1) \cong G_2 = (V_2, E_2)$ if there is a **bijective function** $f: V_1 \rightarrow V_2$ such that for all $(u, v) \in E_1$:


$$(u, v) \in E_1 \text{ iff } (f(u), f(v)) \in E_2$$
- It is **edge-preserving vertex matching**
 - If there is **an edge in the original graph**, there is **an edge after the mapping**; **vice versa**.

Graph Isomorphism Example



$G_1 = (V_1, E_1)$ $G_2 = (V_2, E_2)$
 $V_1 = \{a, b, c, d\}$ $V_2 = \{1, 2, 3, 4\}$
 $E_1 = \{(a,b), (a,c), (a,d), (b,d), (c,d)\}$
 $E_2 = \{(1,3), (1,4), (2,3), (2,4), (3,4)\}$


$(u, v) \in E_1 \text{ iff } (f(u), f(v)) \in E_2$



- $(a,b) \Leftrightarrow (f(a), f(b)) = (3,1)$
- $(a,c) \Leftrightarrow (f(a), f(c)) = (3,2)$
- $(a,d) \Leftrightarrow (f(a), f(d)) = (3,4)$
- $(b,d) \Leftrightarrow (f(b), f(d)) = (1,4)$
- $(c,d) \Leftrightarrow (f(c), f(d)) = (2,4)$


Graph Isomorphism Self-complementary

- A graph G is called **self-complementary** if $G \cong \bar{G}$
 - C_5 and \bar{C}_5 are self-complementary ($C_5 \cong \bar{C}_5$)



Graph Isomorphism

- Showing Isomorphism is not easy
 - No general method which is more efficient than trying all possibilities



Graph Isomorphism

- Showing non-isomorphic is simpler
 - Violate any isomorphic-preserving property
 - Example: Are they isomorphic? NO

